

# Physics Review:

## ② H

- ③ The nucleus is composed of protons and neutrons only
- The protons repel each other electrically
  - The repulsion INCREASES as the protons get closer together
  - In a nucleus the protons are very very very close together
  - To get the protons close enough to be bound by the neutrons requires EXTREME temperature and pressure
  - Those conditions only exist inside of stars and within early universe.

## ④ C, D, F

- A. Find the change in mass or MASS DEFECT.

$$\Delta m = 100 \text{ kg} - 99.87 \text{ kg} = 0.13 \text{ kg}$$


- Use  $E = mc^2$  to find  $E$

$$E = 0.13 \text{ kg} (3 \times 10^8 \text{ m/s})^2 = 1.17 \times 10^{16} \text{ J}$$

B. This is essentially a unit conversion question.

Given: 1 L gasoline =  $3.2 \times 10^7 \text{ J}$

$$1.17 \times 10^{16} \text{ J} \times \left( \frac{1 \text{ L gasoline}}{3.2 \times 10^7 \text{ J}} \right) = 3.65625 \times 10^8 \text{ L gasoline}$$




A. MASS DEFECT:

$$69.938062 \text{ mg} - 69.937704 \text{ mg} = 0.000358 \text{ mg}$$

B. Convert to kg

$$0.000358 \text{ mg} \times \left( \frac{1 \text{ g}}{1000 \text{ mg}} \right) \times \left( \frac{1 \text{ kg}}{1000 \text{ g}} \right) = 3.58 \times 10^{-10} \text{ kg}$$

C.  $\Sigma = mc^2 = 3.58 \times 10^{-10} \text{ kg} (3.0 \times 10^8 \text{ m/s})^2 = 3.222 \times 10^{-7} \text{ J}$










D. MASS DEFECT:

$$226.025402 \text{ u} - 226.020173 \text{ u} = 0.005229 \text{ u}$$

E. CONVERT TO KG:  $0.005229 \text{ u} \times \left( \frac{1.66 \times 10^{-27}}{1 \text{ u}} \right) = 8.68014 \times 10^{-30} \text{ kg}$

F.  $\Sigma = mc^2 = 8.68014 \times 10^{-30} \text{ kg} (3.00 \times 10^8 \text{ m/s})^2 = 7.812 \times 10^{-13} \text{ J}$

(8)  $\Sigma = mc^2 \Rightarrow m = \frac{\Sigma}{c^2} = \frac{4.0 \times 10^{-3} \text{ J}}{(3.00 \times 10^8 \text{ m/s})^2} = 0.0004 \text{ kg} = 0.444 \text{ g}$



(10) The nuclear fusion of Hydrogen into Helium.

(11) A.  $^{212}_{84}\text{Po}$  : 84 protons, 128 neutrons

B.  $^{140}_{58}\text{Ce}$  : 58 protons, 82 neutrons

C.  $^{37}_{17}\text{Cl}$  : 17 protons, 20 neutrons

D.  $^{45}_{21}\text{Sc}$  : 21 protons, 24 neutrons.


E.  $^{80}_{35}\text{Br}$  : 35 protons, 45 neutrons

F.  $^{209}_{83}\text{Bi}$  : 83 protons, 126 neutrons

12. A.  $^{32}_{15}\text{P}$       B.  $^{172}_{70}\text{Yb}$       C.  $^{16}_8\text{O}$       D.  $^{140}_{59}\text{Pr}$

13.  $^{100}_{41}\text{Nb}$

14.  $^{25}_{12}\text{Mg}$



16. See #3. To create the temperature/pressure necessary would require more energy than the reaction would output. Thus it is unfeasible from an economic and energy standpoint.